

*What Is Claimed Is:*

1. An isolated nucleic acid molecule comprising a polynucleotide having a nucleotide sequence at least 90% or more identical to a sequence selected from the group consisting of:
  - (a) a polynucleotide fragment of SEQ ID NO:1 or a polynucleotide fragment of the cDNA sequence included in ATCC Deposit No: 209641 or 209691;
  - (b) a polynucleotide encoding a polypeptide fragment of SEQ ID NO:2 or the cDNA sequence included in ATCC Deposit No: 209641 or 209691;
  - (c) a polynucleotide encoding a polypeptide domain of SEQ ID NO:2 or the cDNA sequence included in ATCC Deposit No: 209641 or 209691;
  - (d) a polynucleotide encoding a polypeptide epitope of SEQ ID NO:2 or the cDNA sequence included in ATCC Deposit No: 209641 or 209691;
  - (e) a polynucleotide encoding a polypeptide of SEQ ID NO:2 or the cDNA sequence included in ATCC Deposit No: 209641 or 209691 having biological activity;
  - (f) a polynucleotide which is a variant of SEQ ID NO:1;
  - (g) a polynucleotide which is an allelic variant of SEQ ID NO:1;
  - (h) a polynucleotide which encodes a species homologue of the SEQ ID NO:2;and
  - (i) a polynucleotide capable of hybridizing under stringent conditions to any one of the polynucleotides specified in (a)-(h), wherein said polynucleotide does not hybridize under stringent conditions to a nucleic acid molecule having a nucleotide sequence of only A residues or of only T residues.
2. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises a nucleotide sequence encoding a mature form or a secreted protein.
3. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises a nucleotide sequence encoding the sequence identified as SEQ ID NO:2 or the coding sequence included in ATCC Deposit No: 209641 or 209691.

4. The isolated nucleic acid molecule of claim 1, wherein the polynucleotide fragment comprises the entire nucleotide sequence of SEQ ID NO:1 or the cDNA sequence included in ATCC Deposit No: 209641 or 209691.
5. The isolated nucleic acid molecule of claim 2, wherein the nucleotide sequence comprises sequential nucleotide deletions from either the C-terminus or the N-terminus.
6. The isolated nucleic acid molecule of claim 3, wherein the nucleotide sequence comprises sequential nucleotide deletions from either the C-terminus or the N-terminus.
7. A recombinant vector comprising the isolated nucleic acid molecule of claim 1.
8. A method of making a recombinant host cell comprising the isolated nucleic acid molecule of claim 1.
9. A recombinant host cell produced by the method of claim 9.
10. The recombinant host cell of claim 9 comprising vector sequences.
11. An isolated polypeptide comprising an amino acid sequence at least 90% or more identical to a sequence selected from the group consisting of:
  - (a) a polypeptide fragment of SEQ ID NO:2 or the encoded sequence included in ATCC Deposit No: 209641 or 209691;
  - (b) a polypeptide fragment of SEQ ID NO:2 or the encoded sequence included in ATCC Deposit No: 209641 or 209691 having biological activity;
  - (c) a polypeptide domain of SEQ ID NO:2 or the encoded sequence included in ATCC Deposit No: 209641 or 209691;
  - (d) a polypeptide epitope of SEQ ID NO:2 or the encoded sequence included in ATCC Deposit No: 209641 or 209691;
  - (e) a mature form of a secreted protein;
  - (f) a full length secreted protein;
  - (g) a variant of SEQ ID NO:2;

- (h) an allelic variant of SEQ ID NO:2; and
- (i) a species homologue of the SEQ ID NO:2.

12. The isolated polypeptide of claim 11, wherein the mature form or the full length secreted protein comprises sequential amino acid deletions from either the C-terminus or the N-terminus.
13. An isolated antibody that binds specifically to the isolated polypeptide of claim 11.
14. A recombinant host cell that expresses the isolated polypeptide of claim 11.
15. A method of making an isolated polypeptide comprising:
  - (a) culturing the recombinant host cell of claim 14 under conditions such that said polypeptide is expressed; and
  - (b) recovering said polypeptide.
16. The polypeptide produced by claim 15.
17. A method for preventing, treating, or ameliorating a medical condition which comprises administering to a mammalian subject a therapeutically effective amount of the polypeptide of claim 11.
18. A method of diagnosing a pathological condition or a susceptibility to a pathological condition in a subject related to expression or activity of a secreted protein comprising:
  - (a) determining the presence or absence of a mutation in the polynucleotide of claim 1;
  - (b) diagnosing a pathological condition or a susceptibility to a pathological condition based on the presence or absence of said mutation.
19. A method of diagnosing a pathological condition or a susceptibility to a pathological condition in a subject related to expression or activity of a secreted protein comprising:

- (a) determining the presence or amount of expression of the polypeptide of claim 11 in a biological sample;
  - (b) diagnosing a pathological condition or a susceptibility to a pathological condition based on the presence or amount of expression of the polypeptide.
20. A method for identifying binding partner to the polypeptide of claim 11 comprising:
- (a) contacting the polypeptide of claim 11 with a binding partner; and
  - (b) determining whether the binding partner effects an activity of the polypeptide.
21. The gene corresponding to the cDNA sequence of SEQ ID NO:2.
22. A method of identifying an activity in a biological assay, wherein the method comprises:
- (a) expressing SEQ ID NO:1 in a cell;
  - (b) isolating the supernatant;
  - (c) detecting an activity in a biological assay; and
  - (d) identifying the protein in the supernatant having the activity.
23. The product produced by the method of claim 22
24. A multimer comprising the polypeptide of claim 11.